

## HYPNOSIS TO MANAGE DISTRESS RELATED TO MEDICAL PROCEDURES: A META-ANALYSIS

Julie B. Schnur, Ilana Kafer, Carolyn Marcus and Guy H. Montgomery

*Mount Sinai School of Medicine, New York, USA*

---

### Abstract

This meta-analysis evaluates the effect of hypnosis in reducing emotional distress associated with medical procedures. PsycINFO and PubMed were searched from their inception through February 2008. Randomized controlled trials of hypnosis interventions, administered in the context of clinical medical procedures, with a distress outcome, were included in the meta-analysis (26 of 61 papers initially reviewed). Information on sample size, study methodology, participant age and outcomes were abstracted independently by 2 authors using a standardized form. Disagreements were resolved by consensus. Effects from the 26 trials were based on 2342 participants. Results indicated an overall large effect size (ES) of 0.88 (95% CI = 0.57–1.19) in favour of hypnosis. Effect sizes differed significantly ( $p < 0.01$ ) according to age (children benefitted to a greater extent than adults) and method of hypnosis delivery, but did not differ based on the control condition used (standard care vs. attention control). Copyright © 2008 British Society of Experimental & Clinical Hypnosis. Published by John Wiley & Sons, Ltd.

---

**Key words:** hypnosis, emotional distress, medical, meta-analysis

### Introduction

*I learned a long time ago that minor surgery is when they do the operation on someone else, not you. Bill Walton (Elston, 2005: 14)*

As the above quotation reveals, medical procedures are often associated with a great deal of emotional distress for patients, and the possible sources of such distress are manifold. The procedures themselves can involve ‘immobilization, darkness (in the case of image-guided procedures), masked strangers, fear for life and health, uncertainty of outcome, and abdication of control’ (Flory, Salazar and Lang, 2007: 304). For those procedures involving anesthesia, patients may possess additional concerns. Research indicates that up to 54.5% of patients fear being unable to ‘wake up’ after surgery, and up to 54% fear ‘waking up’ during surgery (Klafta and Roizen, 1996). Furthermore, 16% fear pain associated with procedures, and on a more general level, up to 62% fear anesthesia (Klafta and Roizen, 1996). Depending on the procedure, there can also be tremendous concerns about post-procedure functioning, including concerns about subsequent pain, impaired daily living, body image (e.g. scarring; Ersek and Denton, 1986; Chun and Velanovich, 2002), and diagnosis/prognosis (i.e. what the surgeon will find (Schnur, Montgomery, Hallquist, Goldfarb, Silverstein and Weltz, 2008b)).

Emotional distress related to medical procedures not only causes direct suffering, but across procedures, pre-procedure emotional stress and distress have been related to a

variety of adverse post-procedure outcomes including: postoperative emotional distress (Munafò and Stevenson, 2001); pain (e.g. (Kain, Servarino, Aleander, Pincus and Mayes, 2000; Cohen, Fouladi and Katz, 2005; Carr, Brockbank, Allen and Strike, 2006; Hong, Jee and Luthardt, 2005; Munafò and Stephenson, 2001)); postsurgery nausea and fatigue (Montgomery and Bovbjerg, 2004); failure to return to work and failure to report improvements in pain and functional ability (Trief, Grant and Fredrickson, 2000); recovery room and total analgesic requirements (Pan, Coghill, Houle, Seid, Lindel and Parker, 2006); dose/demand ratio for patient controlled analgesia (PCA), degree of dissatisfaction with PCA (Ozalp, Sarioglu, Tuncel, Aslan and Kadiogullari, 2003); increased propofol requirements (Osborn and Sandler, 2004; Hong et al., 2005); and worsened wound healing (Ginandes, Brooks, Sando, Jones and Aker, 2003).

Therefore, to reduce both the direct negative experience of emotional distress related to medical procedures, as well as the possible negative downstream consequences of such distress, interventions which significantly reduce distress are needed to improve patient experience. Although pharmacologic interventions can be beneficial, they are not without cost. They can cause their own side effects (Klein, 1991; Hollenhorst, Munte, Friedrich, Heine, Leuwer and Becker, 2001; Flory et al., 2007), and they can be a drain on medical staff time because patients taking sedatives can require increased monitoring and nursing care (Murphy and Brunberg, 1997). An additional issue is that it can be difficult to precisely time when to administer anxiolytic medication; if given too late, such medication may not become effective until after the procedure has already begun (Quirk, Letendre, Ciottono and Lingley, 1989; Murphy and Brunberg, 1997). All of these factors suggest that a non-pharmacologic adjuvant treatment for patient emotional distress would be an important tool for patient care.

Hypnosis is a non-pharmacologic intervention, with no known specific side effects (Rhue, Lynn and Kirsch, 1993; Lynn, Martin and Frauman, 1996), which has been shown in both narrative reviews (Redd, Montgomery and DuHamel, 2001; Flory et al., 2007) and meta-analysis (Montgomery, David, Winkel, Silverstein and Bovbjerg, 2002) to be beneficial in reducing distress related to medical procedures. However, the existing reviews in this area have been limited in several ways. The Redd et al. paper (2001) served to illustrate the effectiveness of hypnosis in the context of medical procedures, but was focused exclusively on the cancer setting, and thus did not include the broader gamut of medical procedures. The paper by Flory and colleagues was a narrative review of a select number of trials, which provided an illustration of the effectiveness of hypnosis for medical procedures, rather than providing a comprehensive review of the literature. Furthermore, neither the Redd et al. nor the Flory et al. papers provided an empirical summary of the data reviewed. The meta-analysis (Montgomery et al., 2002) did provide such empirical data, but was limited in that it focused exclusively on the surgical setting, and neglected individuals undergoing other types of medical procedures (e.g. lumbar puncture, radiotherapy). In addition, the meta-analysis did not limit its inclusion criteria to randomized controlled trials, which are considered the 'gold standard' research design.

Therefore the current literature is lacking a meta-analysis of randomized controlled trials on the use of hypnosis to reduce emotional distress related to a broad range of medical procedures, both surgical and non-surgical. The primary goal of the present study was to remedy this gap in the literature by using meta-analytic techniques with data from published randomized controlled trials to estimate the effectiveness of hypnosis in managing distress related to medical procedures.

## Method

### *Search strategy*

Two electronic databases, PsycInfo and PubMed were searched from their respective inception through the end of February 2008.

For PsycInfo, the major search terms were (mj = hypnosis OR mj = hypnotherapy) AND (mj = distress or mj = medical treatment general or mj = treatment effectiveness evaluation or mj = postsurgical complications or mj = anxiety or mj = stress-management or mj = stress or mj = treatment outcomes or mj = fear or mj = emotional states or mj = relaxation), and the limitations placed on the search were ((DT:PSYI = CHAPTER) or (DT:PSYI = JOURNAL-ARTICLE)) and (LA:PSYI = ENGLISH) and ((MD:PSYI = EMPIRICAL-STUDY) or (MD:PSYI = FIELD-STUDY) or (MD:PSYI = FOLLOWUP-STUDY) or (MD:PSYI = LONGITUDINAL-STUDY) or (MD:PSYI = META-ANALYSIS) or (MD:PSYI = PROSPECTIVE-STUDY) or (MD:PSYI = QUANTITATIVE-STUDY) or (MD:PSYI = RETROSPECTIVE-STUDY) or (MD:PSYI = TREATMENT-OUTCOME-CLINICAL-TRIAL)) and ((PT:PSYI = JOURNAL) or (PT:PSYI = PEER-REVIEWED-JOURNAL) or (PT:PSYI = PEER-REVIEWED-STATUS-UNKNOWN)). This yielded a total of 248 abstracts/titles.

For PubMed, the search terms were: ('Hypnosis'[Mesh] OR 'Hypnosis, Dental'[Mesh])AND('Stress Disorders, Traumatic'[Mesh] OR 'Postoperative Complications'[Mesh] OR 'Treatment Outcome'[Mesh] OR 'Stress, Psychological'[Mesh] OR 'Preoperative Care'[Mesh] OR 'Anxiety'[Mesh] OR 'Depression'[Mesh] OR 'Fear'[Mesh] OR 'Affect'[Mesh] OR 'Relaxation'[Mesh] OR 'Emotions'[Mesh] OR 'Psychology'[Mesh]). The only limitation placed on the search was that the articles had to be written in English. This yielded a total of 1661 abstracts/titles.

### *Selection strategy*

The abstracts/titles of all articles identified by electronic searches (1909 in total) were carefully screened by three of the authors in the study to determine if the abstracts met the following inclusion criteria: (a) published in a peer reviewed journal; (b) full abstract available online; (c) randomized trial; (d) written in English; (e) included at least one control condition; (f) hypnosis was listed as at least one of the intervention conditions; (g) the use of hypnosis was specifically related to a medical or dental procedure (other than childbirth, which was considered to be a phenomenon rather than a procedure); (h) some measure of distress (e.g. anxiety) or emotional well-being (e.g. relaxation) was identified as an outcome variable; (i) the study was not a secondary analysis of an existing data set; (j) the study had sufficient data to calculate an effect size; and (k) the study was not a duplicate (i.e. if an article was cited in both PubMed and PsycInfo, it was only used once).


Subsequent to abstract review, 61 manuscripts were obtained and read in full, independently by two co-authors, each of whom completed a standardized form assessing the inclusion criteria. All four co-authors then met to review these 61 manuscripts. Based on consensus review by all authors, 26 of those 61 papers were included in the meta-analysis. It is interesting to note that although the electronic search was for hypnosis interventions, six papers were retrieved which referred to their interventions as suggestion rather than hypnosis. These papers were included as they met all remaining inclusion criteria, and would provide an interesting comparison to interventions labelled hypnosis. The reasons for exclusion were as follows: in 14 studies, participants were not randomly assigned to study group, 5 studies did not have sufficient data to calculate an effect size,

3 studies were secondary data analyses, 3 studies were not RCTs (i.e. they were review papers), 3 studies did not concern a specific medical procedure, 3 did not have a control condition, in 2 distress was not an outcome, and in one the intervention was not referred to as hypnosis or suggestion. Additionally, one paper was excluded (Lee, 2003) as the intervention was multimodal (hypnosis in addition to cognitive behavioural counselling), rather than being hypnosis focused (see QUORUM Flow Chart, Figure 1).

### Data abstraction and study characteristics

For each of the 26 papers accepted for inclusion into the meta-analysis, relevant data was abstracted using a standardized worksheet. Each paper was abstracted independently by two of the co-authors. Any discrepancies were discussed among the authors with reference to the original manuscript until consensus was reached. Specific data collected included:

- (1) between-group results on distress outcomes needed to calculate an effect size (e.g. means, standard deviation, sample sizes, test statistics);
- (2) how the hypnosis intervention was labelled by the authors, coded as either ‘hypnosis’ or ‘suggestion.’ ‘Hypnosis’ was defined as the actual word hypnosis, as well as its derivatives (e.g. hypnotherapeutic ego strengthening). ‘Suggestion’ was defined as any derivative of the term suggestion (e.g. therapeutic suggestions). It should be noted that although our original intent was to restrict the sample to studies which explicitly labelled their intervention ‘hypnosis,’ we realized upon reviewing the literature retrieved that there were a small but substantial number of studies which called their intervention ‘suggestions’. The methodological descriptions of these ‘suggestion’ interventions were remarkably similar to those for ‘hypnosis’ interven-



**Figure 1.** QUORUM flow chart.

- tions, with the exception that the authors did not label 'suggestion' interventions as 'hypnosis'. We therefore decided to code this data, in the service of conducting an exploratory moderation analysis of whether the intervention label was associated with intervention effect size;
- (3) average participant age, which was examined categorically. Child was defined as less than or equal to age 18;
  - (4) control condition, which was coded as either 'standard care' or as 'attention control.' 'Attention control' was defined as any control condition other than standard care (e.g. blank tape, white noise);
  - (5) method of hypnosis delivery, which was coded as either 'live' or 'recorded.' 'Live' was defined as an intervention with any live, in-person component. 'Recorded' was defined as an intervention with *no* live, in-person component;
  - (6) timing of intervention delivery, which was coded as either 'pre,' 'during,' or 'pre and during.' 'Pre' was defined as an intervention delivered prior to a medical procedure, 'during' was defined as an intervention delivered during the medical procedure with *no* pre-procedure delivery, and interventions which incorporated both delivery approaches were labelled 'pre and during';
  - (7) whether or not the effect size needed to be imputed. This was coded as 'Imputed,' which indicates a paper where means and standard deviations were not provided in text or table format, and where effect sizes had to be derived from test or descriptive statistics (based on Smith, Glass and Miller, 1980).

### *Quantitative data synthesis*

For every study, we calculated an effect size (ES) for each comparison between the intervention of interest (i.e. hypnosis, suggestions) and a control condition on a distress outcome. If a given study had more than one hypnosis condition (e.g. direct hypnosis, indirect hypnosis) or more than one control condition (e.g. attention control, standard care) then an ES was computed for each combination of hypnosis condition and control condition. If a given study had multiple measurements of emotional distress (e.g. at different post-intervention time points, or different outcome measures) an effect size was calculated for each. Effect sizes were not calculated for time points prior to the administration of the intervention.

The effect sizes were calculated using the meta-analysis program by Schwarzer (Schwarzer, 2008) (using the 'Effect Sizes *d*' utility). The effect sizes generated from this utility are *g*, where  $g = (M_c - M_e)/SD$ . In this equation,  $M_c$  represents the mean of the control group on the outcome variable,  $M_e$  represents the mean of the experimental group on the outcome variable, and *SD* represents the pooled standard deviation of the two groups. If a given study did not present means and standard deviations, then the ES was calculated based on test or descriptive statistics following methods suggested by Smith and colleagues (Smith et al., 1980). For ease of interpretation, effect sizes are positive if the hypnosis intervention had the desired beneficial outcome (e.g. less distress or more relaxation), and negative if the hypnosis intervention had the opposite of the desired effect (e.g. more distress or less relaxation).

Once the initial set of effect sizes was generated, a mean effect size for each comparison, for each study, was calculated. This was done by taking the mean of all of the relevant effects for that comparison (i.e. across distress outcome variables).

The effect sizes were then averaged using the meta-analysis program by Schwarzer (2008). The results produced include not only an unweighted overall effect size, but also

an effect size weighted by sample size produced using a random effects model (delta). Both statistics will be reported, since the random effects model does not assume that the set of effect sizes is homogeneous, and we expected the set of effects to display heterogeneity. Additionally, it has been recommended to use the random effects model, as this model is more generalizable to the broader population of studies (Rosenthal and DiMatteo, 2001).

The results yielded by the programme also address two important concerns: the homogeneity of the set of effects, and the 'file-drawer' problem. Homogeneity of the set of effects is critical, as it indicates the trustworthiness of the overall effect size generated by the meta-analysis (Schwarzer, 2008). To examine homogeneity of the effect sizes, we will examine the percentage of variance attributable to sampling error versus the percentage of variance attributable to systematic factors, as well as the  $Q$  statistic. The  $Q$  statistic indicates whether the variability present in the group of effect sizes is significantly greater than chance, and thus suggestive of the presence of potential moderators of intervention effects. To examine publication bias, otherwise known as the 'file-drawer problem,' we used Orwin's (1983) method. Moderator analyses following a finding of heterogeneity of effect sizes (a significant  $Q$  value), were conducted by analyses of variance.

## Results

### *Description of trial, sample and intervention characteristics*

The 26 randomized trials meeting the inclusion criteria yielded 36 effect sizes. Table 1 presents the chief characteristics and effect sizes for each trial. It should be noted that these are  $g$  values, not Cohen's  $d$  values, and therefore may not be identical to the published  $d$  values in the original manuscripts.

The trials were published between 1984 and 2008. The age range of participants was from 4.8 years to 70.3 years. Sample size ranged from 20 to 200. As can be noted from Table 1, the intervention timing and delivery varied across studies. Outcomes assessed in the studies included anxiety (17 studies), general distress or mood disturbance (6 studies), depression (5 studies), behaviour disorders/distress behaviours (4 studies), relaxation (3 studies), tension (2 studies), anger (1 study), fear (1 study), crying (1 study), how traumatic the procedure was considered (1 study) and nervousness (1 study).

### *Quantitative data synthesis: effectiveness of hypnosis compared with control conditions*

The aggregated effect sizes demonstrate that hypnosis had a significant, large and beneficial effect on emotional distress related to medical procedures. Using a random effects model, we found a mean effect size of 0.88 ( $SE = 0.16$ , 95%  $CI = 0.57$  to 1.19), and this value was significantly greater than zero ( $z = 5.58$ ,  $p < 0.0001$ ).

As noted above, the effect sizes used for this analysis were all drawn from published, peer-reviewed journals, which may raise concerns about publication bias. To address this concern, we calculated the number of studies with an effect size of zero required to decrease the overall mean effect size to small or medium (based on Orwin, 1983), and created a funnel plot (Figure 2). Results indicated that 28 studies with effect sizes of zero would be needed to reduce the large intervention effect size of 0.88 found here to a medium effect size of 0.50, and that 123 studies with effect sizes of zero would be needed to reduce the present effect size of 0.88 to a small effect size of 0.20.

**Table 1.** Study characteristics and mean effect sizes (Hedges' *g*)


Study	Procedure	Method <sup>a</sup> /Timing <sup>b</sup> of Hypnosis		Intervention Label <sup>f</sup>	Type of Control Condition <sup>d</sup>	<i>n</i>	Children vs. Adults <sup>e</sup>	ES ( <i>g</i> )
		Intervention	Control					
Ashton et al. 1995	Coronary artery bypass	Live/Pre	Hypnosis	Standard care	22	Adults	0.46	
Blankfield et al. 1995	Coronary artery bypass	Recorded/During	Suggestion	Attention control	63	Adults	0.16	
Block et al. 1991	Mixed surgery	Recorded/During	Suggestion	Attention control	199	Adults	0.04	
Butler et al. 2005	Voiding cystourethrography	Live/Pre and During	Hypnosis	Standard care	44	Children	0.44	
Calipel et al. 2005	Lower abdominal surgery	Live/Pre	Hypnosis	Standard care	50	Children	0.65	
Cruise et al. 1997	Cataract surgery	Recorded/During	Suggestion	Attention control	60	Adults	0.30	
	Cataract surgery	Recorded/During	Suggestion	Attention control	59	Adults	0.17	
de Klerk et al. 2004	Coronary artery bypass	Live/Pre	Hypnosis	Standard care	50	Adults	0.93	
Faymonville et al. 1997	Plastic surgery	Live/During	Suggestion	Standard care	56	Adults	0.05	
Goldmann et al. 1988	Gynecologic surgery	Live/Pre	Hypnosis	Standard care	52	Adults	0.92	
Harandi et al. 2004	Physiotherapy with burn patients	Live/Pre	Hypnosis	Standard care	44	Adults and Children	2.38	
Katcher et al. 1984	Oral surgery	Live/Pre	Hypnosis	Attention control	32	Adults	0.68	
Lang et al. 2000	Interventional radiology	Live/During	Hypnosis	Standard care	161	Adults	0.37	
	Interventional radiology	Live/During	Hypnosis	Attention control	162	Adults	0.24	
Lang et al. 2006	Large core breast biopsy	Live/During	Hypnosis	Standard care	154	Adults	0.37	
	Large core breast biopsy	Live/During	Hypnosis	Attention control	160	Adults	0.19	
Liossi and Hatira 1999	Bone marrow aspirations	Live/Pre and During	Hypnosis	Standard care	20	Children	1.29	
Liossi and Hatira 2003	Lumbar puncture	Live/Pre and During	Hypnosis (Direct)	Standard care	40	Children	2.64	
	Lumbar puncture	Live/Pre and During	Hypnosis (Direct)	Attention control	40	Children	2.11	
	Lumbar puncture	Live/Pre and During	Hypnosis (Indirect)	Standard care	40	Children	2.66	
	Lumbar puncture	Live/Pre and During	Hypnosis (Indirect)	Attention control	40	Children	2.12	

Table 1. Continued

Study	Procedure	Method <sup>a</sup> /Timing <sup>b</sup> of Hypnosis Intervention	Intervention Label <sup>c</sup>	Type of Control Condition <sup>d</sup>	n	Children vs. Adults <sup>e</sup>	ES (g)
Lioosi et al. 2006	Lumbar puncture	Live Pre and During	Hypnosis	Attention control	30	Children	3.14
Liu et al. 1992	Lumbar puncture	Live/Pre and During	Hypnosis	Standard care	30	Children	3.16
	Total abdominal hysterectomy	Recorded/During	Suggestion	Attention control	49	Adults	0
Marc et al. 2007	Total abdominal hysterectomy	Recorded/During	Suggestion	Attention control	50	Adults	0
	Surgical abortion	Live/Pre and During	Hypnosis	Standard care	29	Adults	0.17
Massarini et al. 2005	Mixed surgery	Live/Pre	Hypnosis	Standard care	42	Adults and Children	0.69
Montgomery et al. 2002	Excisional breast biopsy	Live/Pre	Hypnosis	Standard care	20	Adults	1.25
Montgomery et al. 2007	Breast conserving surgery	Live/Pre	Hypnosis	Attention control	200	Adults	1.11
Saadat et al. 2006	Ambulatory surgery	Live/Pre	Hypnosis	Standard care	50	Adults	2.33
	Ambulatory surgery	Live/Pre	Hypnosis	Attention control	52	Adults	1.69
Schnur et al. 2008a	Excisional breast biopsy	Live/Pre	Hypnosis	Attention control	90	Adults	0.79
Stalpers et al. 2005	Mixed cancer radiotherapy	Live/Pre and During	Hypnosis	Standard care	69	Adults	0
	Hysterectomy, myomectomy, or GYN laparotomy	Recorded/Pre	Suggestion	Attention control	40	Adults	0.52
Wood et al. 1990	Tonsillectomy and adenoidectomy	Recorded/During	Suggestion	Attention control	49	Children	0.12
	Tonsillectomy and adenoidectomy	Recorded/During	Suggestion	Attention control	50	Children	0.36

Note: a) 'live' was defined as an intervention with a live, in-person component, 'recorded' was defined as an intervention with *no* live, in-person component; b) 'pre' was defined as an intervention delivered prior to a medical procedure, 'during' was defined as an intervention delivered during the medical procedure with *no* pre-procedure delivery, and interventions which incorporated both delivery approaches were labeled 'pre and during'; c) 'hypnosis' was defined as the authors using the actual word hypnosis, as well as its derivatives (e.g. hypnotherapeutic ego strengthening), to describe the intervention whereas suggestion was defined as the authors using any derivative of the term suggestion (e.g. therapeutic suggestions); d) 'attention control' was defined as any control condition other than standard care (e.g. blank tape, white noise); e) Child was defined as <age 18.





**Figure 2.** Effect size by study sample size (funnel plot).

Results of homogeneity tests indicated that the sample of effect sizes was heterogeneous ( $Q = 231.31$ ,  $df = 35$ ,  $p < 0.0001$ ). Only 12.86% of the variance was explained by sampling error, indicating that 87.14% of the variance was explained by systematic factors. This suggests that moderators are likely to be present.

#### *Impact of moderators on hypnosis effects*

Based on the degree of heterogeneity in the data set, we proceeded to test potential moderator variables. Table 2 presents the descriptive and inferential statistics for these tests. Results indicate that: a) hypnosis is significantly more effective for children than for adults ( $p < 0.001$ ); b) sample size is significantly and inversely correlated with effect size ( $p < 0.05$ ); c) average effect size does not differ significantly whether hypnosis is compared to an attention control or to a standard care control condition ( $p > 0.23$ ); d) interventions were significantly more effective when labelled 'hypnosis' than when labelled 'suggestion' ( $p < 0.002$ ); e) hypnosis was significantly more effective when it was delivered at least in part using a 'live' administration method as compared to when the intervention was delivered via audio recording ( $p < 0.005$ ); f) hypnosis was significantly more effective when it was delivered at least in part prior to the medical procedure (our Pre and Pre and During conditions) rather than solely during the medical procedure (post-hoc Tukey  $p < 0.05$ ) (see Table 2); and g) effect sizes did not differ significantly between those studies where effect sizes had to be imputed and those studies where effect sizes did not have to be imputed ( $p > 0.08$ ). As can be seen in Table 1, there is a great deal of overlap between the study characteristics of live vs. recorded, the timing of the intervention delivery, and hypnosis label. Studies which were delivered 'live' also tended to be labelled 'hypnosis' and to be delivered prior to the medical procedure, while those which were delivered via recording were more likely to be delivered during the procedure and to be labelled 'suggestion'. The overlapping nature of these factors prevented an analysis of the effects of the moderators using more complex statistical models (e.g. two- and three-way interactions). With regard to the finding that sample size was inversely correlated with effect size, a funnel plot (Figure 2) revealed that this correlation may be driven by one larger sample size study with an effect size of zero (Block, Ghoneim, Sum Ping and Ali, 1991). The correlation between sample size and effect size with this particular study removed was non-significant ( $p = 0.08$ ).

**Table 2.** Effect sizes and significance tests for moderation effects

Moderator of interest	$n_{\text{effects}}$	Mean ( <i>SD</i> ) ES by class	Test statistic	<i>p</i>
Child vs. adult <sup>a</sup>	Child = 11 Adult = 23	Child = 1.70 (1.16) Adult = .55 (.60)	$F(1,33) = 14.62$	0.0006
Sample size (continuous)	32		$r = -.34$	0.0438
Control condition (standard care vs. attention control) <sup>b</sup>	Standard Care = 18 Attention Control = 18	Standard Care = 1.15 (1.02) Attention Control = 0.76 (0.91)	$F(1,35) = 1.46$	0.2352
Label (hypnosis vs. suggestions) <sup>c</sup>	Hypnosis = 26 Suggestion = 10	Hypnosis = 1.26 (0.99) Suggestion = 0.17 (0.17)	$F(1,35) = 11.79$	0.0016
Method of hypnosis delivery (live vs. recorded) <sup>d</sup>	Live = 27 Recorded = 9	Live = 1.22 (1.0) Recorded = 0.19 (0.18)	$F(1,35) = 9.34$	0.0043
Hypnosis administration time (pre vs. during vs. pre and during) <sup>e</sup>	Pre = 13 During = 13 Pre and During = 10	Pre = 1.11 (0.64) During = 0.18 (0.14) Pre and During = 1.77 (1.22)	$F(1,35) = 13.17$	<0.0001
Imputation <sup>f</sup>	Imputed = 14 Not imputed = 22	Imputed = 0.61 (.70) Not imputed = 1.18 (1.07)	$F(1,35) = 3.09$	0.0878

*Note:* a) Child was defined as <age 18; b) 'attention control' was defined as any control condition other than standard care (e.g. blank tape, white noise); c) 'hypnosis' was defined as the actual word hypnosis, as well as its derivatives (e.g. hypnotherapeutic ego strengthening), whereas suggestion was defined as any derivative of the term suggestion (e.g. therapeutic suggestions); d) 'live' was defined as an intervention with a live, in-person component, 'recorded' was defined as an intervention with no live, in-person component; e) 'pre' was defined as an intervention delivered prior to a medical procedure, 'during' was defined as an intervention delivered during the medical procedure with no pre-procedure delivery, and interventions which incorporated both delivery approaches were labeled 'pre and during'; f) Tukey post-hoc tests revealed that mean During effect size was significantly lower than mean effect sizes for Pre or Pre and During ( $p < 0.05$ ). No other comparisons were significantly different ( $p > 0.05$ ); g) 'Imputed' indicates a paper where means and standard deviations were not provided in text or table format, and where effect sizes had to be derived (based on Smith et al., 1980).

## Discussion

This meta-analysis represents the most extensive review of randomized trials to date on the effects of hypnosis to reduce emotional distress related to medical procedures. The results indicate that approximately 82% of patients undergoing medical procedures who receive hypnosis exhibit lower levels of emotional distress relative to patients in a control condition. The finding of a large effect size in favour of hypnosis supports its more widespread dissemination.

The overall effectiveness of hypnosis for controlling emotional distress associated with medical procedures is consistent with a previous meta-analysis on the effects of hypnosis on negative affect in surgical settings (Montgomery, Weltz, Seltz and Bovbjerg, 2002), which found a hypnosis effect size of 1.07. Effect sizes in the previous meta-analysis may have been larger as the analysis included non-randomized studies and excluded interventions not explicitly labelled as hypnosis. Both the present meta-analysis and the previous one resulted in effect sizes for hypnosis in the large range. The consistency of these findings suggests that hypnosis can improve the patient experience related to medical procedures.

Examination of moderating factors yielded several interesting findings. First, the finding that hypnosis has a medium effect size for adults and a large effect size for children supports the utility of hypnosis regardless of patient age. The finding of a significantly larger clinical effect for children than for adults may be explained by research indicating that children may be more responsive to suggestion (Morgan and Hilgard, 1972). However, it should also be noted that some of the hypnosis interventions delivered to children in the present set of studies tended to be more involved (e.g. more treatment components – self- and hetero-hypnosis, greater number of or longer intervention sessions) than those delivered to adults. Therefore, it is difficult to state with certainty whether the moderation effect is explainable by age differences or by intervention dose differences. Future studies could disentangle this finding by having similarly dosed interventions delivered to both children and adults. Second, the finding that hypnosis appears equally effective whether compared to an attention control group or to a standard care control group suggests that the effects of hypnosis are not merely due to attention. Third, the results indicated that the most effective hypnosis interventions were those that used the label ‘hypnosis’. However, these interventions also tended to be delivered both live and prior to medical procedures, making the exact determination of unique moderating effects problematic.

Future research manipulating the label of the hypnosis intervention, the timing of the intervention, and the presentation (live vs. recorded) of the intervention would be helpful in identifying the most effective hypnosis delivery method for medical procedures. From a practical perspective, explicitly labelling interventions as hypnosis has been found to increase clinical effect sizes (Schoenberger, Kirsch, Gearan, Montgomery and Pastyrnak, 1997). Using the label ‘hypnosis’ may be an efficient clinical approach for increasing patient benefit. Lastly, the results indicated that studies with larger sample sizes tended to have smaller effect sizes, due in part to a few small studies with extremely large effect sizes, and one particular large sample size study with an effect size of zero. One might speculate that there were a greater number of smaller sample size studies with larger ESs, as authors may have stopped their trials once beneficial effects were clearly evident.

Table 1 can be used to clarify clinical research areas which have received greater attention to date (e.g. coronary artery bypass surgery), and more importantly, those

which remain in need of further empirical study. For example, no clinical trials were identified that focused on the use of hypnosis to reduce distress related to non-invasive imaging procedures (e.g. MRI, CT scan) or to relatively newer procedures (e.g. stereotactic radiosurgery, brachytherapy). Similarly, only one study focused on radiotherapy, one of the three standard oncology treatments. Finally, the vast majority of the studies assessed the reduction of negative emotions (a critical outcome), whereas few assessed any positive emotions which might result from hypnosis (e.g. calm, relaxation, joy, self-confidence). As a result, the literature may be telling only half the story of the benefits of hypnosis for medical procedures.

One limitation of this meta-analysis is that we were unable to address the question of the potential moderating effects of hypnotic suggestibility, as too few papers reported these data. Although an important consideration, the overall effect size documented here suggests that the vast majority of patients can benefit from a hypnosis intervention to reduce distress related to medical procedures, regardless of level of hypnotic suggestibility.

In summary, the data strongly support the use of hypnosis as a non-pharmacologic intervention to reduce emotional distress associated with medical procedures, and suggest that the more widespread adoption of hypnosis could improve the quality of life of millions of patients undergoing medical procedures.

## **Acknowledgements**

Preparation of this manuscript was supported by the American Cancer Society (RSGPB-04-213-01-CPPB, PF-05-098-01-CPPB) and the National Cancer Institute (1R25CA129094-01, K07 CA131473-01).

## **References**

- Ashton RC Jr, Whitworth GC, Seldomridge JA, Shapiro PA, Michler RE, Smith CR et al. (1995) The effects of self-hypnosis on quality of life following coronary artery bypass surgery: preliminary results of a prospective, randomized trial. *Journal of Alternative and Complementary Medicine* 1: 285–90.
- Blankfield RP, Zyzanski SJ, Flocke SA, Alemagno S, Scheurman K (1995) Taped therapeutic suggestions and taped music as adjuncts in the care of coronary-artery-bypass patients. *American Journal of Clinical Hypnosis* 37: 32–43.
- Block RI, Ghoneim MM, Sum Ping ST, Ali MA (1991) Efficacy of therapeutic suggestions for improved postoperative recovery presented during general anesthesia. *Anesthesiology* 75: 746–55.
- Butler LD, Symons BK, Henderson SL, Shortliffe LD, Spiegel D (2005) Hypnosis reduces distress and duration of an invasive medical procedure for children. *Pediatrics* 115: e77–e85.
- Calipel S, Lucas-Polomeni MM, Wodey E, Ecoffey C (2005) Premedication in children: hypnosis versus midazolam. *Paediatric Anaesthesia* 15: 275–81.
- Carr E, Brockbank K, Allen S, Strike P (2006) Patterns and frequency of anxiety in women undergoing gynaecological surgery. *Journal of Clinical Nursing* 15: 341–52.
- Chun K, Velanovich V (2002) Patient-perceived cosmesis and satisfaction after breast biopsy: comparison of stereotactic incisional, excisional, and wire-localized biopsy techniques. *Surgery* 131: 497–501.
- Cohen L, Fouladi RT, Katz J (2005) Preoperative coping strategies and distress predict postoperative pain and morphine consumption in women undergoing abdominal gynecologic surgery. *Journal of Psychosomatic Research* 58: 201–9.
- Cruise CJ, Chung F, Yogendran S, Little D (1997) Music increases satisfaction in elderly outpatients undergoing cataract surgery. *Canadian Journal of Anaesthesia* 44: 43–8.

- de Klerk JE, du Plessis WF, Steyn HS, Botha M (2004) Hypnotherapeutic ego strengthening with male South African coronary artery bypass patients. *American Journal of Clinical Hypnosis* 47: 79–92.
- Elston G (2005) *The Wild World of Sports*. Bloomington, Indiana: AuthorHouse.
- Ersek RA, Denton DR (1986) Breast biopsy technique: a plea for cosmesis. *The Southern Medical Journal* 79: 167–70.
- Faymonville ME, Mambourg PH, Joris J, Vrijens B, Fissette J, Albert A et al. (1997) Psychological approaches during conscious sedation. Hypnosis versus stress reducing strategies: a prospective randomized study. *Pain* 73: 361–7.
- Flory N, Salazar GM, Lang EV (2007) Hypnosis for acute distress management during medical procedures. *International Journal of Clinical and Experimental Hypnosis* 55: 303–17.
- Ginandes C, Brooks P, Sando W, Jones C, Aker J (2003) Can medical hypnosis accelerate post-surgical wound healing? Results of a clinical trial. *American Journal of Clinical Hypnosis* 45: 333–51.
- Goldmann L, Ogg TW, Levey AB (1988) Hypnosis and daycase anaesthesia. A study to reduce pre-operative anxiety and intra-operative anaesthetic requirements. *Anaesthesia* 43: 466–9.
- Harandi AA, Esfandani A, Shakibaei F (2004) The effect of hypnotherapy on procedural pain and state anxiety related to physiotherapy in women hospitalized in a burn unit. *Contemporary Hypnosis* 21: 28–34.
- Hollenhorst J, Munte S, Friedrich L, Heine J, Leuwer M, Becker H et al. (2001) Using intranasal midazolam spray to prevent claustrophobia induced by MR imaging. *American Journal of Roentgenology* 176: 865–8.
- Hong JY, Jee YS, Luthardt FW (2005) Comparison of conscious sedation for oocyte retrieval between low-anxiety and high-anxiety patients. *Journal of Clinical Anesthesia* 17: 549–53.
- Kain ZN, Servarino F, Aleander GM, Pincus S, Mayes LC (2000) Preoperative anxiety and postoperative pain in women undergoing hysterectomy. A repeated-measures design. *Journal of Psychosomatic Research* 49: 417–22.
- Katcher A, Segal H, Beck A (1984) Comparison of contemplation and hypnosis for the reduction of anxiety and discomfort during dental surgery. *American Journal of Clinical Hypnosis* 27: 14–21.
- Klafta JM, Roizen MF (1996) Current understanding of patients' attitudes toward and preparation for anesthesia: a review. *Anesthesia and Analgesia* 83: 1314–21.
- Klein DS (1991) Prevention of claustrophobia induced by MR imaging: use of alprazolam. *American Journal of Roentgenology* 156: 633.
- Lang EV, Benotsch EG, Fick LJ, Lutgendorf S, Berbaum ML, Berbaum KS et al. (2000) Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomised trial. *Lancet* 355: 1486–90.
- Lang EV, Berbaum KS, Faintuch S, Hatsiopolou O, Halsey N, Li X et al. (2006) Adjunctive self-hypnotic relaxation for outpatient medical procedures: a prospective randomized trial with women undergoing large core breast biopsy. *Pain* 126: 155–64.
- Lioffi C, Hatira P (1999) Clinical hypnosis versus cognitive behavioral training for panic management with pediatric cancer patients undergoing bone marrow aspirations. *International Journal of Clinical and Experimental Hypnosis* 47: 104–16.
- Lioffi C, Hatira P (2003) Clinical hypnosis in the alleviation of procedure-related pain in pediatric oncology patients. *International Journal of Clinical and Experimental Hypnosis* 51: 4–28.
- Lioffi C, White P, Hatira P (2006) Randomized clinical trial of local anesthetic versus a combination of local anesthetic with self-hypnosis in the management of pediatric procedure-related pain. *Health Psychology* 25: 307–15.
- Liu WH, Standen PJ, Aitkenhead AR (1992) Therapeutic suggestions during general anaesthesia in patients undergoing hysterectomy. *British Journal of Anaesthesia* 68: 277–81.
- Lynn SJ, Martin DJ, Frauman DC (1996) Does hypnosis pose special risks for negative effects? *International Journal of Clinical and Experimental Hypnosis* 44: 7–19.

- Marc I, Rainville P, Verreault R, Vaillancourt L, Masse B, Dodin S (2007) The use of hypnosis to improve pain management during voluntary interruption of pregnancy: an open randomized preliminary study. *Contraception* 75: 52–8.
- Massarini M, Rovetto F, Taglaferri C, Preoperative Hypnosis study group (2005) Preoperative hypnosis. A controlled study to assess the effects on anxiety and pain in the postoperative period. *European Journal of Clinical Hypnosis* 6: 8–15.
- Montgomery GH, Bovbjerg DH (2004) Pre-surgery distress and specific response expectancies predict post-surgery outcomes in surgery patients confronting breast cancer. *Health Psychology* 23: 381–7.
- Montgomery GH, Bovbjerg DH, Schnur JB, David D, Goldfarb A, Weltz CR et al. (2007) A randomized clinical trial of a brief hypnosis intervention to control side effects in breast surgery patients. *Journal of the National Cancer Institute* 99: 1304–12.
- Montgomery GH, David D, Winkel G, Silverstein JH, Bovbjerg DH (2002) The effectiveness of adjunctive hypnosis with surgical patients: a meta-analysis. *Anesthesia and Analgesia* 94: 1639–45.
- Montgomery GH, Weltz CR, Seltz G, Bovbjerg DH (2002) Brief pre-surgery hypnosis reduces distress and pain in excisional breast biopsy patients. *International Journal of Clinical and Experimental Hypnosis* 50: 17–32.
- Morgan AH, Hilgard ER (1972) Age differences in susceptibility to hypnosis. *International Journal of Clinical and Experimental Hypnosis* 21: 78–85.
- Munafo MR, Stevenson J (2001) Anxiety and surgical recovery. Reinterpreting the literature. *Journal of Psychosomatic Research* 51: 589–96.
- Murphy KJ, Brunberg JA (1997) Adult claustrophobia, anxiety and sedation in MRI. *Magnetic Resonance Imaging* 15: 51–4.
- Orwin RG (1983) A fail safe N for effect size in meta-analysis. *Journal for Educational Statistics* 8: 157–9.
- Osborn TM, Sandler NA (2004) The effects of preoperative anxiety on intravenous sedation. *Anesthesia Progress* 51: 46–51.
- Ozalp G, Sarioglu R, Tuncel G, Aslan K, Kadiogullari N (2003) Preoperative emotional states in patients with breast cancer and postoperative pain. *Acta Anaesthesiologica Scandinavica* 47: 26–9.
- Pan PH, Coghill R, Houle TT, Seid MH, Lindel WM, Parker RL et al. (2006) Multifactorial preoperative predictors for postcesarean section pain and analgesic requirement. *Anesthesiology* 104: 417–25.
- Quirk ME, Letendre AJ, Ciottoni RA, Lingley JF (1989) Anxiety in patients undergoing MR imaging. *Radiology* 170: 463–6.
- Redd WH, Montgomery GH, DuHamel KN (2001) Behavioral intervention for cancer treatment side-effects. *Journal of National Cancer Institute* 93: 810–23.
- Rhue JW, Lynn SJ, Kirsch I (1993) *Handbook of Clinical Hypnosis*. Washington, DC: American Psychological Association.
- Rosenthal R, DiMatteo MR (2001) Meta-analysis: recent developments in quantitative methods for literature reviews. *Annual Review of Psychology* 52: 59–82.
- Saadat H, Drummond-Lewis J, Maranets I, Kaplan D, Saadat A, Wang SM et al. (2006) Hypnosis reduces preoperative anxiety in adult patients. *Anesthesia and Analgesia* 102: 1394–6.
- SAS Institute I (2002) *Statistical Analysis Software (SAS)* [Computer software]. Cary, N.C.
- Schnur JB, Bovbjerg DH, David D, Tatrow K, Goldfarb AB, Silverstein JH et al. (2008a) Hypnosis decreases presurgical distress in excisional breast biopsy patients. *Anesthesia and Analgesia* 106: 440–4.
- Schnur JB, Montgomery GH, Hallquist MN, Goldfarb AB, Silverstein JH, Weltz CR et al. (2008b) Anticipatory psychological distress in women scheduled for diagnostic and curative breast cancer surgery. *International Journal of Behavioral Medicine* 15: 21–8.

- Schoenberger NE, Kirsch I, Gearan P, Montgomery GH, Pastyrnak SL (1997) Hypnotic enhancement of a cognitive behavioral treatment for public speaking anxiety. *Behavior Therapy* 28: 127–40.
- Schwarzer R (2008) *Statistics Software for Meta-Analysis* [Computer software].
- Smith ML, Glass GV, Miller TI (1980) *The Benefits of Psychotherapy*. Baltimore: Johns Hopkins University Press.
- Stalpers LJ, da Costa HC, Merbis MA, Fortuin AA, Muller MJ, van Dam FS (2005) Hypnotherapy in radiotherapy patients: a randomized trial. *International Journal of Radiation Oncology, Biology, Physics* 61: 499–506.
- Trief PM, Grant W, Fredrickson B (2000) A prospective study of psychological predictors of lumbar surgery outcome. *Spine* 25: 2616–21.
- van der Laan WH, van Leeuwen BL, Sebel PS, Winograd E, Baumann P, Bonke B (1996) Therapeutic suggestion has not effect on postoperative morphine requirements. *Anesthesia and Analgesia* 82: 148–52.
- Wood WE, Gibson W, Longo D (1990) Moderation of morbidity following tonsillectomy and adenoidectomy: a study of awareness under anesthesia. *International Journal of Pediatric Otorhinolaryngology* 20: 93–105.

*Address for correspondence:*

*Dr Julie B. Schnur*

Department of Oncological Sciences  
Box 1130, Mount Sinai School of Medicine  
1425 Madison Avenue  
New York, NY 10029-6574  
Phone: 212-659-5644  
Fax: 212-849-2564  
Email: julie.schnur@mssm.edu

Copyright of Contemporary Hypnosis is the property of John Wiley & Sons, Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.